

Fine-structure constant variability: surprises for laboratory atomic spectroscopy and cosmological evolution of quasar spectra

Jacob D. Bekenstein*

*Racah Institute of Physics, Hebrew University of Jerusalem
Givat Ram, Jerusalem 91904 ISRAEL*

(Dated: February 2, 2008)

Calculation of the Dirac hydrogen atom spectrum in the framework of dynamical fine structure constant (α) variability discloses a small departure in the laboratory from Sommerfeld's formula for the fine structure shifts, possibly measurable today. And for a distant object in the universe, the wavelength shift of a spectral line specifically ascribable to cosmological α variation is found to depend differently on the quantum numbers than in the conventional view. This last result clashes with the conventional wisdom that an atom's spectrum can change with cosmological time only through evolution of the α parameter in the energy eigenvalue formula, and thus impacts on the Webb group's analysis of fine structure intervals in quasar absorption lines (which has been claimed to disclose cosmological α evolution). In particular, analyzing together a mix of quasar absorption lines from different fine structure multiplets can bias estimates of cosmological α variability.

PACS numbers: 98.80.Es, 06.20.Jr, 98.62.Ra, 32.30.Jc, 71.15.Rf

Early theoretical speculations about temporal variability of the fine-structure constant α [1, 2, 3] (an excellent review is provided by Uzan [4]) have assumed urgency in view of the claimed cosmological variability in fine-structure multiplets seen in absorption in distant quasar spectra [5, 6]. In the Webb group's analysis it is assumed that the spectrum of an atom in an intergalactic cloud at redshift z can be gotten from its laboratory spectrum by merely replacing the laboratory value of α by that at redshift z . This assumption is crucial to the group's strategy for reducing experimental errors by analyzing simultaneously lines from different fine-structure multiplets of different elements.

The mentioned assumption takes it for granted that in a world where α varies, the *form* of the Dirac hamiltonian (the most direct tool for determining the fine structure splittings) relevant for an electron in an atom is the same as in standard physics. Is such an "electromagnetic equivalence principle" correct?

In one paradigm cosmological α variability [7] occurs in discrete jumps, each precipitated by a phase transition in which spontaneous symmetry breaking changes a certain vacuum expectation value, and thereby the value of α . In this point of view it seems reasonable for the Dirac Hamiltonian to retain its textbook form, at least during each cosmological interval of constant α .

Things are different in the much earlier paradigm, first contemplated by Jordan [1] and Dicke [3], where α is a dynamical field. Both workers emphasized that in a covariant theory, temporal dynamical α variability must be accompanied by spatial variability. Concerns, first articulated by Dicke [3], that spatial variability necessarily entails observable violations of the weak equivalence prin-

ciple have proved groundless, at least within the general framework of dynamical α variability proposed by us [8]: its violations of the equivalence principle lie beyond foreseeable experimental sensitivities [9].

As we show here for the hydrogen atom, dynamical α variability according to the framework modifies the form of the Dirac hamiltonian. One consequence is a small deviation of the hydrogen atom's spectrum in today's laboratory from that given by Sommerfeld's formula [Eq. (15) below]. Dynamical α variability also implies nontrivial evolution of atomic spectra with cosmological epoch (time). Estimation of α 's cosmological evolution from quasar fine-structure multiplets of mixed provenance must take cognizance of this effect.

In its most convenient form, the framework for α variability is a theory of *constant* charges where the permittivity and the reciprocal permeability of the vacuum both equal $e^{-2\psi}$, where ψ is a scalar field [9, 10]. One consequence is that the coupling "constant" α of Coulomb's law, which is inversely proportional to the vacuum's permittivity, varies as $e^{2\psi}$. Another is that the electromagnetic field tensor $f^{\mu\nu}$ and its dual $*f^{\mu\nu}$ obey the equations

$$*f^{\mu\nu}{}_{;\nu} = 0, \quad (1)$$

$$(e^{-2\psi} f^{\mu\nu})_{;\nu} = \frac{4\pi}{c} j^\mu \quad (2)$$

with j^μ the electric current. The coupling between $e^{-2\psi}$ and $f^{\mu\nu}$ induces an electromagnetic contribution to the source term of the scalar field equation,

$$\psi_{,;\mu}{}^{,\mu} = 4\pi\kappa^2\sigma - \frac{\kappa^2}{2}e^{-2\psi}f_{\mu\nu}f^{\mu\nu}, \quad (3)$$

where the density σ is associated with particles and κ is the parameter of the theory with dimensions of reciprocal energy times charge,. Whenever j^μ can be described as a sum of contributions from discrete particles, σ is obtained by replacing in j^μ the product of each particle's

*Electronic address: bekenste@vms.huji.ac.il;
URL: <http://www.phys.huji.ac.il/~bekenste/>

charge and the corresponding 4-velocity by the derivative $\partial mc^2/\partial\psi$ of the particle's mass m . So in this theory particle mass is, in principle, a function of the ambient ψ field.

In flat spacetime the *physical* static solution of Eqs. (1)-(3) for a pointlike charge e_0 in terms of its formal Coulomb potential $\Phi = e_0/r$ and its true electric field, $\mathbf{E} \equiv c\{f^{01}, f^{02}, f^{03}\}$, is [9]

$$\mathbf{E} = -e^{2\psi}\nabla\Phi = -\nabla(\kappa^{-1}\tan(\kappa\Phi + \chi)) \quad (4)$$

$$e^\psi = \sec(\kappa\Phi + \chi), \quad (5)$$

with χ a constant of integration. A second such constant has been set to unity by redefinition of the charge [9]. This restricts our description to epochs for which $e^\psi > 1$. Now χ is evidently equivalent to $\exp(\psi_{\text{cosm}})$, the cosmological value of e^ψ at the relevant epoch, which plays the role of asymptotic value of $e^{\psi(r)}$. Solution (4)-(5) thus depends on an *adiabatically* varying parameter. We assume that $|\chi| \ll 1$ throughout the epochs of interest.

Solution (4)-(5) applies as well to multiple pointlike particles at rest [9] with Φ replaced by the Coulomb potential coming from all the charges. According to Ref. 9, such a solution is consistent arbitrarily near a source only if the latter's mass m is subject to the condition $\kappa^2\partial mc^2/\partial\psi = -e_0 d\psi/d\Phi$ with both sides *evaluated* at the relevant particle's position. The theory thus fixes each m 's dependence on ψ . By Eq. (5) the condition is

$$\partial mc^2/\partial\Phi = -e_0 \tan^2(\kappa\Phi + \chi) \quad (6)$$

The derivative here gives the change of m reflecting a local change of the total Coulomb potential Φ caused by charges elsewhere (the charge's self-potential, Φ_S , cannot change, of course; since the framework does not allow truly point charges, $\Phi_S < \infty$ [9]).

The integral of Eq. (6) is

$$mc^2 = C - e_0\kappa^{-1}\tan(\kappa\Phi + \chi) + e_0\Phi \quad (7)$$

with C an integration constant. In terms of the free particle's rest mass m_0 ($m \rightarrow m_0$ for $\Phi \rightarrow \Phi_S$) at the epoch in question, $C = m_0 + e_0\kappa^{-1}\tan(\kappa\Phi_S + \chi) - e_0\Phi_S$. Now from Ref. 9 it follows that κ^{-1} is no more than an order of magnitude below Planck's energy divided by the elementary charge. Thus, for example, even at distance 10^{-17} cm from an electron (if such accuracy in position has any meaning in light of the *Zitterbewegung*), $e_0\Phi$ receives a contribution 18 orders *below* Planck energy and $\kappa|\Phi|$ and $\kappa|\Phi_S|$ are both smaller than 10^{-18} . Since we have no reasons to expect that χ is this tiny at a generic epoch, we may regard both $\kappa|\Phi|$ and $\kappa|\Phi_S|$ as small compared to χ . Thus for a (pointlike) electron with charge e_0 immersed in the Coulomb potential Φ_N of a nucleus at a cosmological epoch specified by χ , we obtain by expanding Eq. (7) to first order in $\kappa\Phi_N = \kappa(\Phi - \Phi_S)$, and neglecting in the tangent's argument $\kappa\Phi_S$ as compared to χ that

$$m = m_0 + \delta m \approx m_0 - e_0 c^{-2} \tan^2 \chi \cdot \Phi_N. \quad (8)$$

Consider now the Dirac hamiltonian $\hat{H}$ for the electron in an electric field $\mathbf{E}$. In an inertial frame

$$\hat{H} = (-i\hbar c\boldsymbol{\alpha} \cdot \nabla + (m_0 + \delta m)c^2\beta + e_0 UI) \quad (9)$$

with I the 4×4 unit matrix and β and the triplet $\boldsymbol{\alpha}$ the four familiar Dirac matrices [12]. By minimal coupling the potential U in which the electron moves must be such that $\mathbf{E} = -\nabla U$. From Eq. (4) we infer that

$$U = \kappa^{-1}\tan(\kappa\Phi + \chi) - \kappa^{-1}\tan(\kappa\Phi_S + \chi), \quad (10)$$

at the electron's position, the added constant ensuring that U vanishes asymptotically for any χ , just as does Φ_N . We include the electron's Φ_S in Φ because of nonlinearity of the theory [see Eqs. (4)-(5)]. Expanding U to $\mathcal{O}(\Phi_N)$ and again dropping $\kappa\Phi_S$ in comparison with χ , we get

$$U \approx \sec^2 \chi \cdot \Phi_N. \quad (11)$$

With help of Eqs. (8) and (11), Eq. (9) may be rewritten as

$$\hat{H} = \hat{H}_0 + \delta\hat{H} \quad (12)$$

$$\hat{H}_0 = (-i\hbar c\boldsymbol{\alpha} \cdot \nabla + m_0 c^2\beta + e_0\Phi_N I) \quad (13)$$

$$\delta\hat{H} = (I - \beta)e_0 \tan^2 \chi \cdot \Phi_N \quad (14)$$

Now the mass of the nucleus (proton) is also subject to the same "correction" as in Eq. (8): the proton's charge is $-e_0$ but it feels the *electron's* potential $-\Phi_N$. However, we do not include this mass "correction" in $\hat{H}$ just as we do not add an extra $(-e_0)(-\Phi_N)$ on account of the nucleus to the *potential* term in Dirac's equation. We likewise drop from $\hat{H}$ the free proton's mass (analog of m_0) since it cannot affect transition wavelengths.

Eq. (13) is the textbook Dirac hamiltonian for an electron (charge e_0) in the external Coulomb potential Φ_N . Let $\alpha_* \equiv e_0^2/\hbar c$ (to be distinguished from the coupling "constant" α). In terms of the principal quantum number $n = 1, 2, \dots$ and the total angular momentum quantum number $j = \frac{1}{2}, \frac{3}{2}, \dots$, its eigenvalues, to $\mathcal{O}(\alpha_*^4)$, are [11]

$$E_0 = m_0 c^2 \left[1 - \frac{\alpha_*^2}{2n^2} - \frac{\alpha_*^4}{2n^3} \left(\frac{1}{j + \frac{1}{2}} - \frac{3}{4n} \right) \right]. \quad (15)$$

The term of $\mathcal{O}(\alpha_*^2)$ is the Bohr energy level; that of $\mathcal{O}(\alpha_*^4)$ is Sommerfeld's fine structure correction.

In Eq. (12) $\delta\hat{H}$ represents the perturbation coming from cosmological α evolution. We shall now show that it modifies the conventional expression for the wavelength λ of the transition $n \rightarrow n - \Delta n$, $j \rightarrow j - \Delta j$ to

$$\frac{1}{\lambda} = \frac{m_0 c \alpha_*^2}{4\pi\hbar} \times \left[\Delta \frac{1}{n^2} + \alpha^2 \Delta \left(\frac{1}{n^3(j + \frac{1}{2})} - \frac{3}{4n^4} \right) + \frac{1}{2} \alpha (\alpha - \alpha_*) \Delta \frac{1}{n^4} \right]. \quad (16)$$

This may be contrasted with the conventional one-parameter formula coming just from Eq. (15), namely Eq. (16) with $\alpha_* \mapsto \alpha$ everywhere.

If Ξ denotes the unit-normalized eigenspinor of the eigenvalue problem $\hat{H}_0\Xi = E_0\Xi$, then to first order in perturbation theory the cosmological evolution shift in the energy level is

$$\delta E = \mathcal{I} \tan^2 \chi \quad (17)$$

$$\mathcal{I} \equiv \int \Xi^\dagger (I - \beta) \Xi e_0 \Phi_n d^3x \quad (18)$$

What is $\tan^2 \chi$ in Eq. (17)? Since α/α_* is the reciprocal permittivity of the vacuum, and $\exp(2\psi_{\text{cosm}}) = \sec^2 \chi$, we may write

$$\tan^2 \chi = \sec^2 \chi - 1 = (\alpha - \alpha_*)/\alpha_* \quad (19)$$

Thus the permittivity is unity when $\chi = 0$. The precise epoch in question depends on the cosmological model; we defer consideration of this issue to a future publication. Our choice precludes study of epochs with $\alpha < \alpha_*$.

To calculate $\mathcal{I}$ one first notes that $I - \beta$ is the matrix $\text{diag}(0, 0, 2, 2)$ so that with Ξ written as the bispinor $\begin{pmatrix} \xi \\ \eta \end{pmatrix}$,

$$\mathcal{I} = 2 \int \eta^\dagger \eta e_0 \Phi_n d^3x. \quad (20)$$

Now in the usual representation of α in terms of the Pauli matrices σ , $\hat{H}_0\Xi = E_0\Xi$ takes the form [12]

$$-i\hbar c \sigma \cdot \nabla \eta + m_0 c^2 \xi + e_0 \Phi_n \xi = E_0 \xi, \quad (21)$$

$$-i\hbar c \sigma \cdot \nabla \xi - m_0 c^2 \eta + e_0 \Phi_n \eta = E_0 \eta. \quad (22)$$

By requiring, in spherical polar coordinates $\{r, \theta, \varphi\}$, that Ξ be an eigenfunction of the z -component as well as of the square of the total angular momentum, $\hat{\mathbf{J}}$ (eigenvalues $m\hbar$ and $j(j+1)\hbar^2$, respectively), one is led to the two possible forms [12]

$$\Xi = \begin{pmatrix} F(r) \mathcal{Y}_{j-1/2}^{jm} \\ -if(r) \mathcal{Y}_{j+1/2}^{jm} \end{pmatrix} \quad \text{or} \quad \Xi = \begin{pmatrix} G(r) \mathcal{Y}_{j+1/2}^{jm} \\ -ig(r) \mathcal{Y}_{j-1/2}^{jm} \end{pmatrix} \quad (23)$$

with $\mathcal{Y}_\ell^{jm}(\theta, \varphi)$ a spinorial spherical harmonic. The two forms differ in their parity for given j ; the first corresponds to the case $\ell = j - \frac{1}{2}$ in nonrelativistic theory of the hydrogen atom, while the second has $\ell = j + \frac{1}{2}$. If the spinorial harmonics are normalized according to $\int \mathcal{Y}_\ell^{jm\dagger} \mathcal{Y}_\ell^{jm} \sin \theta d\theta d\varphi = 1$, then the normalization of Ξ requires $\int_0^\infty (F^2 + f^2) r^2 dr = 1$ and similarly for G and g (as will be clear all four functions can be chosen real).

In terms of the dimensionless variables $x = rm_0 c/\hbar$, $\epsilon = \mathcal{E}_0/m_0 c^2$ and $k = j + \frac{1}{2}$, the equations for the first form of Ξ can be cast into the form [12]

$$(\epsilon - 1 + \alpha_*/x)F - [d/dx + (k+1)/x]f = 0, \quad (24)$$

$$(\epsilon + 1 - \alpha_*/x)f + [d/dx + (k-1)/x]F = 0 \quad (25)$$

These admit the series solution

$$F = e^{\sqrt{1-\epsilon^2}x} x^\gamma \sum_{\nu=0} a_\nu x^\nu, \quad (26)$$

$$f = e^{\sqrt{1-\epsilon^2}x} x^\gamma \sum_{\nu=0} b_\nu x^\nu.$$

Consistency between the two relations connecting a_0 and b_0 requires that

$$\gamma = -1 + \sqrt{k^2 - \alpha_*^2} \quad (27)$$

(the negative root is unphysical). In order for the sums in Eqs. (26) to terminate at $\nu = n' = 0, 1, \dots$ (otherwise F and f would diverge), one must have

$$\epsilon = \left[1 + \frac{\alpha_*^2}{[\sqrt{k^2 - \alpha_*^2} + n']^2} \right]^{-1/2}. \quad (28)$$

Expanding this to $\mathcal{O}(\alpha_*^4)$ gives the Sommerfeld spectrum, Eq. (15), with $n \equiv n' + j + \frac{1}{2}$.

Because Φ_n is spherically symmetric, the normalization of the spinorial harmonics allows us to write

$$\mathcal{I} = 2 \int_0^\infty f^2 e_0 \Phi_n r^2 dr, \quad (29)$$

where $\Phi_n = -e_0/r$. If Ξ is not normalized to unity, then in passing to the integration variable x we must write

$$\mathcal{I} = -\frac{2m_0 c^2 \alpha_* \int_0^\infty f^2 x dx}{\int_0^\infty (F^2 + f^2) x^2 dx}. \quad (30)$$

The exact relativistic wavefunctions [11] are rather cumbersome, but are not needed if all we want is to calculate $\mathcal{I}$ to $\mathcal{O}(\alpha_*^4)$. Thus starting with $a_0 = 1$, we used *Mathematica* to work out, for a definite n' , all the other a_ν and b_ν to $\mathcal{O}(\alpha_*^{n'+1})$ by including in the recursion relations [12] expressions for γ and ϵ correct to $\mathcal{O}(\alpha_*^{n'+1})$. The series in F and f [see Eq. (26)] turn out to be of the form $\sum_{i=0} U_i(\alpha_* x) \alpha_*^{2i}$ and $\alpha \sum_{i=0} V_i(\alpha_* x) \alpha_*^{2i}$, where U_i and V_i are polynomials of order n' . Up to normalization, $U_0(\alpha_* x)$ is the associated Laguerre polynomial familiar from nonrelativistic theory of the hydrogen atom. The correctness of the so obtained F up to terms with $i = 1$ and of the $i = 0$ term in f was checked by verifying that the Hellmann-Feynman theorem [12],

$$\frac{\partial E_0}{\partial \alpha_*} = \langle \Xi | \frac{\partial \hat{H}_0}{\partial \alpha_*} | \Xi \rangle = m_0 c^2 \frac{\int_0^\infty (F^2 + f^2) x dx}{\int_0^\infty (F^2 + f^2) x^2 dx}, \quad (31)$$

is satisfied with E_0 given by Eq. (15).

A change in the integration variable in Eq. (30) to $y = \alpha_* x$ (essentially the radial coordinate in units of the Bohr radius) brings out an extra factor α_* in the ratio of integrals. Since f carries an overall factor of α_* , the $\mathcal{O}(\alpha_*^4)$ contribution δE can be computed by including only the $i = 0$ terms of f in the numerator of Eq. (30) and of F in the denominator, and totally discarding f in the denominator. The result of exact integrations is

$$\mathcal{I} = -\frac{m_0 c^2 \alpha_*^4}{n^3} \left(\frac{1}{j + \frac{1}{2}} - \frac{1}{2n} \right) \quad (32)$$

The equations for the second form of Ξ in Eq. (23) are obtained from Eqs. (26)-(28) by the replacements $F \mapsto$

$G, f \mapsto g$ and $k \mapsto -k$. The coefficients a_ν and b_ν are thereby changed, but γ and ϵ are not, except that now $n' = 1, 2, \dots$ (n is still $n' + j + \frac{1}{2}$). Repeating the procedure outlined above we find the analog of $\mathcal{I}$ to have exactly the same form as Eq. (32). One consequence is that, at least to $\mathcal{O}(\alpha_*^4)$, the energies of levels with like j but opposite parity are still degenerate, as in the exact Dirac hydrogen spectrum.

From Eqs. (15), (17), (19) and (32) we deduce formula (16) for the transition wavelength. It shows an unconventional dependence on the cosmological “fossil” α_* , the value of α when the vacuum’s permittivity was unity. In principle fitting a generic collection of laboratory measured transition wavelengths with formula (16) should allow separate determination of α and $\alpha - \alpha_*$. A reliable nonzero result for $\alpha - \alpha_*$ (which if the Webb group is right, might be of order $10^{-5}\alpha$) would establish dynamical α variability according to the theories represented by the framework as opposed to phase transition variability. Possibly evidence for this phenomenon lurks in the residuals of extant measurements of the hydrogen spectrum.

Turning to quasar spectra we remark that *observed* quasar absorption line wavelengths must first be corrected by dividing them by the absorber’s redshift factor $1 + z$. In practice $1 + z$ is taken as the ratio of the observed wavelength of a Bohr transition ($n \neq 0$) to the corresponding laboratory wavelength. The procedure precludes separate determination of the α_* in the prefactor of (16); any difference between it and α is absorbed in the mentioned correction, and α_* and α can only be distinguished by the terms *inside* the square bracket in Eq. (16). The correction also makes it unnecessary to allow for the cosmological evolution of m_0 ; its variation gets absorbed in the mentioned correction.

By subtracting the expression given by Eq. (16) for λ_l^{-1} for the laboratory line from λ_z^{-1} , its analog for the corresponding redshift-corrected quasar line, we get

$$\frac{1}{\lambda_z} - \frac{1}{\lambda_l} = \frac{m_0 c \alpha_l^3 (\alpha_z - \alpha_l)}{2\pi\hbar} \cdot \Delta \left[\frac{1}{n^3(j + \frac{1}{2})} - \frac{1}{2n^4} \right], \quad (33)$$

where we have replaced α_* in the prefactor by α_l in accordance with our earlier remarks, as well as dropped terms of $\mathcal{O}((\alpha_l - \alpha_*)^2)$ and $\mathcal{O}((\alpha_z - \alpha_l)^2)$ in light of the assumption $|\chi| \ll 1$. Contrast Eq. (33) with the conventional formula implicit in the Webb group’s analysis,

$$\frac{1}{\lambda_z} - \frac{1}{\lambda_l} = \frac{m_0 c \alpha_l^3 (\alpha_z - \alpha_l)}{2\pi\hbar} \cdot \Delta \left[\frac{1}{n^3(j + \frac{1}{2})} - \frac{3}{4n^4} \right] \quad (34)$$

We observe that formulae (33) and (34) predict exactly the same $\lambda_z^{-1} - \lambda_l^{-1}$ for different members of a *single* fine structure multiplet (constant n and Δn) because the formulae differ by a term ($\propto n^{-4}$) which is constant within the multiplet. We conclude that comparison of fine structure *intervals* in a quasar with the corresponding ones in the laboratory, with each multiplet taken separately (as done in older work by Webb’s group [5] and its predecessors) cannot distinguish between the α variability from phase transitions and dynamical α variability according to the framework.

However, Webb’s group has recently analyzed jointly lines from different multiplets and in different species [6]. This procedure *is* sensitive to the difference between formulae (33) and (34) and their nonhydrogen counterparts. Whereas the factors in square brackets in the two formulae sometimes have similar values, there are also glaring discrepancies. Often the conventional formula underestimates the quantity in question by a factor of 2, and for the transition $2p_{3/2} \rightarrow 4s_{1/2}$ it underestimates it by a factor of 6. For the transition $2s_{1/2} \rightarrow 3p_{1/2}$ the conventional formula *overestimates* $\lambda_z^{-1} - \lambda_l^{-1}$ by a factor of 31.5 and gets the sign wrong. Clearly if α varies dynamically, a reliable many multiplet estimate of the cosmological α variation from quasar absorption spectra must use formula (33). And consideration of it is indispensable for discriminating between the two mentioned paradigms of α variability.

I thank Jonathan Oppenheim for useful comments and the Israel Science Foundation (grant 129/00) for support.

-
- [1] P. Jordan, *Naturwiss.* **25**, 513 (1937); *Z. f. Physik* **113**, 660 (1939).
 - [2] E. Teller, *Phys. Rev.* **73**, 801 (1948); K. P. Stanyukovich, *Sov. Phys. Doklady* **7**, 1150 (1963); G. Gamow, *Phys. Rev. Letters* **19**, 759, 913 (1967).
 - [3] R. H. Dicke, *Science* **129**, 621 (1959) and *The Theoretical Significance of Experimental Relativity* (Gordon and Breach, New York 1965).
 - [4] J. P. Uzan, e-print hep-ph/0205340.
 - [5] J. K. Webb, V. V. Flambaum, C. W. Churchill, M. J. Drinkwater and J. D. Barrow, *Phys. Rev. Letters* **82**, 884 (1999), and references therein.
 - [6] J. K. Webb, M. T. Murphy, V. V. Flambaum, V. A. Dzuba, J. D. Barrow, C. W. Churchill, J. X. Prochaska and A. M. Wolfe, *Phys. Rev. Letters* **87**, 091301 (2001).
 - [7] Z. Chacko, C. Grojean and M. Perelstein, hep-ph/0204142.
 - [8] J. D. Bekenstein, *Phys. Rev. D* **25**, 1527 (1982).
 - [9] J. D. Bekenstein, *Phys. Rev. D* **66**, 123514 (2002).
 - [10] H. B. Sandvik, J. D. Barrow and J. Magueijo, *Phys. Rev. Lett.* **88**, 031302 (2002).
 - [11] V. B. Berestetskii, E. M. Lifshitz and L. P. Pitaevskii, *Quantum Electrodynamics* (Pergamon, Oxford 1982).
 - [12] E. Merzbacher, *Quantum Mechanics*, Sec. Edition. (Wiley, New York 1970), pp. 442, 603-607.